

5-2004

Computation of the Ramsey Number $R(B_3, K_5)$

Andrey Babak

Rochester Institute of Technology

Stanislaw Radziszowski

Rochester Institute of Technology

Kung-Kuen Tse

Kean University

Follow this and additional works at: <http://scholarworks.rit.edu/article>

Recommended Citation

A. Babak, S.P. Radziszowski, and K.K. Tse, Computation of the Ramsey Number $R(B_3, K_5)$. *Bulletin of the Institute of Combinatorics and its Applications*, 41 (2004), 71-76.

This Article is brought to you for free and open access by RIT Scholar Works. It has been accepted for inclusion in Articles by an authorized administrator of RIT Scholar Works. For more information, please contact ritscholarworks@rit.edu.

Computation of the Ramsey Number $R(B_3, K_5)$

Andrey Babak, and
Stanislaw P. Radziszowski
Department of Computer Science
Rochester Institute of Technology
Rochester, NY 14623
spr@cs.rit.edu

Kung-Kuen Tse
Department of Computer Science
and Mathematics
Kean University
Union, NJ 07083
ktse@kean.edu

Abstract

In 1989, George R. T. Hendry presented a table of two-color graph Ramsey numbers $R(G, H)$ for all pairs of graphs G and H having five vertices, with the exception of seven cases. Until now, only two of these open cases were solved. This work eliminates another one by computing $R(B_3, K_5) = 20$, where $B_3 = K_2 + \overline{K_3}$ is the book graph of order 5. In addition, we show that for these parameters there exists a unique up to isomorphism critical graph. The results are based on computer algorithms. Among the four remaining open cases in Hendry's table, the most notable is that of K_5 versus K_5 , for which it is known that $43 \leq R(K_5, K_5) \leq 49$.

Keywords: Ramsey numbers, graph algorithms

AMS subject classification: 05C55

1. Overview

Let G and H be simple graphs. Graph F will be called a (G, H) -graph if it does not contain a subgraph isomorphic to G , and such that the complement \overline{F} has no subgraph isomorphic to H . A $(G, H; n)$ -graph is a (G, H) -graph of order n . Let $\mathcal{R}(G, H)$ and $\mathcal{R}(G, H; n)$ denote the set of all (G, H) -graphs and $(G, H; n)$ -graphs, respectively. The Ramsey number $R(G, H)$ is defined to be the least $n > 0$ such that there is no $(G, H; n)$ -graph, or equivalently, it is the least positive integer n such that every

2-coloring of the edges of K_n contains a subgraph isomorphic to G in the first color or a subgraph isomorphic to H in the second color.

In the sequel, P_k and C_k denote a simple path and cycle, respectively, on k vertices. Further, $B_k = K_2 + \overline{K}_k$ is the k -page book of order $k + 2$, which can be seen as a graph formed by k triangular pages sharing one common edge. Note that we could define equivalently $B_k = K_1 + K_{1,k}$, where $K_{1,k}$ is a star with k spikes. Finally, let $W_k = K_1 + C_{k-1}$ denote a wheel with $k - 1$ spokes.

In 1989, George R. T. Hendry [1] presented a table of Ramsey numbers $R(G, H)$ for all pairs of graphs G and H having five vertices, with the exception of seven cases. Until now, only two of these open cases have been solved. A regularly updated survey by the second author [4] reports on old and the most recent results on various types of Ramsey numbers, including those of the form $R(G, H)$. In particular, [4] lists the developments related to all seven cases missing in the Hendry's table, and gives references to papers discussing them. For graphs G and H of order less than 5, or G of order at most 4 and H of order 5, all Ramsey numbers $R(G, H)$ have been known since the computation of $R(K_4, K_5) = 25$ [3] in 1995.

In this work, we eliminate one of the open cases by computing $R(B_3, K_5) = 20$. This result improves the bounds $20 \leq R(B_3, K_5) \leq 22$ given in [1]. In addition, we show that for these parameters there exists a unique up to isomorphism critical graph, i.e. $|\mathcal{R}(B_3, K_5; 19)| = 1$. We have also completed a full enumeration of all $(B_3, K_5; 18)$ -graphs, and found that there are exactly 3376 of them. Our results, described in more detail in the next section, are based on computer algorithms.

The remaining open cases of Ramsey numbers for graphs on at most five vertices are: $25 \leq R(K_5 - P_3, K_5) \leq 28$, $27 \leq R(W_5, K_5) \leq 29$, $30 \leq R(K_5 - e, K_5) \leq 34$, and $43 \leq R(K_5, K_5) \leq 49$ (see [4] for references to all bounds). Among them, definitely the most famous one and the most difficult to compute, is that of K_5 versus K_5 . While some very strong evidence has been presented to support the conjecture that $R(K_5, K_5) = 43$, there is little hope for a computational proof of the latter, or even for any improvement of the current lower or upper bound. In contrast, further progress on improving bounds for the other three open cases can be expected in the not so distant future. More specifically, we speculate as follows. We feel that the case of $R(W_5, K_5)$ could be attacked with a method similar to the approach in this paper, though more computational effort would be needed. The evaluation of $R(K_5 - P_3, K_5)$ might be at least as hard to obtain as the result $R(K_4, K_5) = 25$, since $K_5 - P_3$ contains K_4 .

The computation of the exact value of $R(K_5 - e, K_5)$ seems to be still much more difficult, but some improvement of the current bounds (especially upper) should be possible.

2. Enumerations and Results

We will use the same notation as in [5]. If G is a graph, then VG and EG are its vertex set and edge set, respectively. If $v \in VG$, then $N_G(v) = \{w \in VG \mid vw \in EG\}$, and $\deg_G(v) = |N_G(v)|$. The subgraph of G induced by W will be denoted by $G[W]$. Also, for $v \in VG$, define the induced subgraphs $G_v^+ = G[N_G(v)]$ and $G_v^- = G[VG - N_G(v) - \{v\}]$.

Note that if $G \in \mathcal{R}(B_3, K_m; n)$ and $v \in VG$, then necessarily $G_v^+ \in \mathcal{R}(K_{1,3}, K_m; d)$, where $d = \deg_G(v)$, and $G_v^- \in \mathcal{R}(B_3, K_{m-1}; n - d - 1)$. Hence, G_v^+ must be simply a disjoint union of paths and cycles, and G_v^- is of the same type as G , but for $m - 1$.

It is a straightforward exercise to find all 104 graphs in $\mathcal{R}(K_{1,3}, K_5)$, and fairly simple algorithms are sufficient to generate all 65694 graphs in $\mathcal{R}(B_3, K_4)$. The statistics of both families by the number of graphs with fixed number of vertices is given in Table I. Observe that $R(K_{1,3}, K_5) = 13$ and $R(B_3, K_4) = 14$.

s	$ \mathcal{R}(K_{1,3}, K_5; s) $	$ \mathcal{R}(B_3, K_4; s) $
1	1	1
2	2	2
3	4	4
4	7	10
5	10	25
6	16	86
7	19	326
8	21	1518
9	14	7000
10	7	23462
11	2	28838
12	1	4410
13		12
total	104	65694

Table I. Statistics for $\mathcal{R}(K_{1,3}, K_5)$ and $\mathcal{R}(B_3, K_4)$.

Using the graphs reported in Table I as G_v^+ and G_v^- , the family of graphs $\mathcal{R}(B_3, K_5; n)$ can be constructed by applying a gluing algorithm to $G_v^+ \in \mathcal{R}(K_{1,3}, K_5; s)$ and $G_v^- \in \mathcal{R}(B_3, K_4; t)$ for all possible s and t satisfying $s + t + 1 = n$. The gluing algorithm used in this work was simpler than, but similar to, that described in [3, 5]. Also, it had to include some modifications needed in order to avoid the graph B_3 instead of K_4 or C_4 . The computations were completed in three stages as follows.

All $(B_3, K_5; 18)$ -graphs were obtained by performing gluing of graphs G_v^+ to G_v^- as above for $s \in \{4, 5, 6, 7, 8\}$ and $t = 17 - s$. No gluing for $s \geq 9$ needs to be done, since it is easy to see that no $(B_3, K_5; 18)$ -graph can have minimum degree 9 or higher. The statistics of results by the number of edges and the minimum degree is presented in Table II.

All $(B_3, K_5; 19)$ -graphs were obtained in two ways: by performing gluing as above for $s \in \{5, 6, 7, 8\}$, $t = 18 - s$, and independently by constructing and (B_3, K_5) -filtering one-vertex extensions of all 3376 $(B_3, K_5; 18)$ -graphs. Both paths led to the same unique $(B_3, K_5; 19)$ -graph, which is cyclic and regular of degree 6, with the edges connecting pairs of vertices belonging to \mathcal{Z}_{19} in distances 1, 7, and 8.

e	d	4	5	6	7	total
51			2			2
52			6			6
53			42			42
54		1	195	19		215
55		14	448	65		527
56		32	572	208		812
57		19	439	321		779
58		2	246	252		500
59			121	142		263
60			60	82		142
61			19	40		59
62			3	16		19
63				4	6	10
total		68	2153	1149	6	3376

Table II. Statistics for $(B_3, K_5; 18)$ -graphs, $e = |E|$, d is the minimum degree.

Similarly, all $(B_3, K_5; 20)$ -graphs were obtained in two ways: by performing gluing as above for $s \in \{6, 7, 8, 9\}$, $t = 19 - s$, and independently by

constructing and (B_3, K_5) -filtering all one-vertex extensions of the unique $(B_3, K_5; 19)$ -graph. Both paths led to no graphs, and thus $R(B_3, K_5) = 20$.

The graphs $\mathcal{R}(B_3, K_5; 18)$ are really not needed, neither is the uniqueness of the critical graph, to claim that $R(B_3, K_5) = 20$. However, these provided additional strong correctness tests of the enumeration results, since several graphs and graph families were generated more than once by different algorithms.

Theorem. $R(B_3, K_5) = 20$.

Proof. The computations and results described above prove that there does not exist any $(B_3, K_5; 20)$ -graph, so $R(B_3, K_5) \leq 20$. It is easy to verify that a cyclic graph with the edges joining vertices belonging to \mathcal{Z}_{19} , which are in distance 1, 7 or 8, has no B_3 and no $\overline{K_5}$. This implies the lower bound. ■

Three separate implementations of the algorithms were prepared and their results compared. The computations were performed as a part of the MS project by the first author, were verified by the second author, and, independently, all reported graph families were enumerated by the third author. The computational effort of this project was moderate — all computations could now be repeated overnight on a local departmental network (150+ machines). If we were only computing the value of $R(B_3, K_5)$, without completing the exhaustive enumeration of all $(B_3, K_5; \geq 18)$ -graphs, then the whole computation would be still much faster.

A general utility program for graph isomorph rejection, *nauty* [2], together with other graph manipulation tools, written by Brendan McKay, was used extensively. All graphs in $\mathcal{R}(B_3, K_5; 18)$ are available from the second author.

References

- [1] G. R. T. Hendry, Ramsey Numbers for Graphs with Five Vertices, *Journal of Graph Theory*, **13** (1989) 245–248.
- [2] B. D. McKay, nauty users' guide (version 1.5), Technical Report TR-CS-90-02, Computer Science Department, Australian National University, 1990. Source code available at <http://cs.anu.edu.au/people/bdm/nauty>.
- [3] B. D. McKay and S. P. Radziszowski, $R(4, 5) = 25$, *Journal of Graph Theory*, **19** (1995) 309–322.
- [4] S. P. Radziszowski, Small Ramsey Numbers, *Electronic Journal of Combinatorics*, Dynamic Survey 1, revision #9, July 2002, <http://www.combinatorics.org/>.
- [5] S. P. Radziszowski and Kung-Kuen Tse, A Computational Approach for the Ramsey Numbers $R(C_4, K_n)$, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **42** (2002) 195–207.