

Rochester Institute of Technology

RIT Scholar Works

Articles

Faculty & Staff Scholarship

4-2019

Near-Complete Genome Sequences of Vesicular Stomatitis Virus Indiana Laboratory Strains HR and T1026R1 and Plaque Isolates 22-20 and 22-25

Thomas M. Russell

Rochester Institute of Technology

Evan E. Santo

Rochester Institute of Technology

Lisa Golebiewski

Rochester Institute of Technology

Nathan S. Haseley

Illumina, Inc.

Maureen Ferran

Rochester Institute of Technology

Follow this and additional works at: <https://scholarworks.rit.edu/article>



Part of the [Immunology and Infectious Disease Commons](#), and the [Virology Commons](#)

Recommended Citation

n Russell TM, Santo EE, Golebiewski L, Haseley NS, Ferran MC. 2019. Near-complete genome sequences of vesicular stomatitis virus Indiana laboratory strains HR and T1026R1 and plaque isolates 22-20 and 22-25. *Microbiol Resour Announc* 8:e00012-19. <https://doi.org/10.1128/MRA.00012-19>.

This Article is brought to you for free and open access by the Faculty & Staff Scholarship at RIT Scholar Works. It has been accepted for inclusion in Articles by an authorized administrator of RIT Scholar Works. For more information, please contact ritscholarworks@rit.edu.



Near-Complete Genome Sequences of Vesicular Stomatitis Virus Indiana Laboratory Strains HR and T1026R1 and Plaque Isolates 22-20 and 22-25

Thomas M. Russell,^a Evan E. Santo,^{a*} Lisa Golebiewski,^a Nathan S. Haseley,^{a*} Maureen C. Ferran^a

^aThomas H. Gosnell School of Life Sciences, Rochester Institute of Technology, Rochester, New York, USA

ABSTRACT We report four near-complete genome sequences of vesicular stomatitis virus (VSV) Indiana obtained with Sanger and Illumina next-generation sequencing, namely, laboratory strains HR (heat resistant) and T1026R1 and isolates 22-20 and 22-25. Previously, only the M gene of these viruses had been sequenced, and these sequences were not deposited in GenBank.

Vesicular stomatitis virus (VSV) is a member of the genus *Vesiculovirus* in the family *Rhabdoviridae* and has an ~11-kb nonsegmented, negative-sense, single-stranded RNA genome that encodes the nucleoprotein (N), matrix (M), glycoprotein (G), large (L), and phosphoprotein (P) proteins (1). We report the near-complete genome sequences of VSV Indiana laboratory strains HR (heat resistant) (2, 3) and T1026R1 (4) and plaque isolates 22-20 and 22-25, derived from Indiana parent strain 22 (03/87-CR-B, number 22), which was originally isolated from an infected cow (5). HR encodes a wild-type (wt) M protein and is therefore a virus that suppresses interferon (IFN) and inhibits host transcription. T1026R1 encodes a M51R mutation in the M protein which abrogates these functions of M (6–8). The M genes of 22-20 and 22-25 were reported to be identical, even though these isolates are discordant for IFN production in chicken embryo cells (9, 10). To determine if another viral component regulates IFN suppression, we sequenced the majority of these four virus genomes.

Virus stocks were propagated in Vero cells at a multiplicity of infection of 0.0001. Once 80 to 90% of the cells exhibited a cytopathic effect, the supernatant was harvested, centrifuged at $4,000 \times g$, aliquoted, and stored at -80°C . At least two virus stocks of each strain were sequenced and found to be identical. Sanger sequencing was performed first, and Illumina sequencing confirmed these results. Sanger and Illumina sequences were aligned for each virus to generate a consensus genome. For Sanger sequencing, the total RNA was extracted from L929 cells at 4 h postinfection (RNAqueous-4-PCR kit, Ambion). Reverse transcription, PCR, product purification, and sequencing were performed as previously described (11). For Illumina MiSeq sequencing, RNA was extracted from 1 ml of virus stock (QIAamp UltraSens virus kit, Qiagen). RNA quality and concentration were determined (Bioanalyzer Agilent RNA 6000 Nano-Assay), and a library was constructed (TruSeq Stranded mRNA sample preparation kit, Illumina). The samples were pooled, and the multiplexed samples produced an average of 207,990 paired-end reads per sample (2×151 read length). Files were demultiplexed (bcl2fastq version 2.19.0), and adapters and low-quality reads were removed (Trimomatic version 0.36) (12). Processed, cleaned reads were mapped to the human reference genome (GRCh38+gencode28) with STAR version 2.6.0c (13). Unmapped reads were then assembled *de novo* with SPAdes version 3.11.1 (14).

Sanger and Illumina contigs were aligned to each other and mapped to the NCBI reference sequence (GenBank accession number [NC_001560](https://www.ncbi.nlm.nih.gov/nuccore/NC_001560)) with CodonCode Aligner (CodonCode Corp.). Open reading frames (ORFs) were predicted with ORF Finder (Web

Citation Russell TM, Santo EE, Golebiewski L, Haseley NS, Ferran MC. 2019. Near-complete genome sequences of vesicular stomatitis virus Indiana laboratory strains HR and T1026R1 and plaque isolates 22-20 and 22-25. *Microbiol Resour Announc* 8:e00012-19. <https://doi.org/10.1128/MRA.00012-19>.

Editor Jelle Matthijnsens, KU Leuven

Copyright © 2019 Russell et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Maureen C. Ferran, mcsfbi@rit.edu.

* Present address: Evan E. Santo, Department of Pathology, Weill Cornell Medical College, New York, New York, USA; Nathan S. Haseley, Clinical Genomics, Illumina, Inc., San Diego, California, USA.

E.E.S., L.G., and N.S.H. contributed equally to this work.

Received 8 January 2019

Accepted 1 March 2019

Published 4 April 2019

version, NCBI) (15). Genome comparison of HR and T1026R1 confirmed the M51R mutation in the T1026R1 M protein and revealed a novel amino acid substitution in the T1026R1 G protein (S431A). Comparison of 22-25 and 22-20 revealed a previously unidentified D-to-G amino acid substitution at position 52 of the 22-20 M protein (D52G). A single amino acid change was identified in the 22-20 M protein, which suggests that this protein alone suppresses the IFN response. The genomes recovered ranged from 11,115 to 11,143 nucleotides in length, obtaining 99.59 to 99.84% of the genomes relative to the reference sequence. The missing nucleotides are located in noncoding regions at the ends of the genomes.

Data availability. Genome sequences of 22-20, 22-25, T1026R21, and HR were deposited in GenBank (accession numbers [MH919396](#), [MH919397](#), [MH919398](#), and [MH919399](#), respectively). Raw Illumina reads were submitted to the Sequence Read Archive (BioProject number [PRJNA508804](#)).

ACKNOWLEDGMENTS

We thank Jean Lucas-Lenard for generously providing the HR and T1026R1 viruses and Luis L. Rodriguez for isolating the parent strain 22 and providing primer sequences. We are also grateful to Philip I. Marcus, who plaque purified and generously provided the 22-20 and 22-25 virus isolates. We also thank the University of Rochester Genomics Research Center (URGC) for the library preparation, next-generation sequencing, data processing through *de novo* assembly, and technical support.

This work was supported by NIH grant R15 A1058969, the Thomas H. Gosnell School of Life Sciences, and the College of Science at Rochester Institute of Technology.

REFERENCES

- Wagner RR, Rose JK. 1996. *Rhabdoviridae*: the viruses and their replication, p 561-573. In Fields BN, Knipe DM, Howley PM (ed), *Fundamental virology*, 3rd ed. Lippincott-Raven, Philadelphia, PA.
- Marcus PI, Sekellick MJ. 1987. Interferon induction by viruses. XV. Biological characteristics of interferon induction-suppressing particles of vesicular stomatitis virus. *J Interferon Res* 7:269-284. <https://doi.org/10.1089/jir.1987.7.269>.
- Wagner RR, Huang AS. 1966. Inhibition of RNA and interferon synthesis in Krebs-2 cells infected with vesicular stomatitis virus. *Virology* 28:1-10. [https://doi.org/10.1016/0042-6822\(66\)90300-X](https://doi.org/10.1016/0042-6822(66)90300-X).
- Stanners CP, Francoeur AM, Lam T. 1977. Analysis of VSV mutant with attenuated cytopathogenicity: mutation in viral function, P, for inhibition of protein synthesis. *Cell* 11:273-281. [https://doi.org/10.1016/0092-8674\(77\)90044-7](https://doi.org/10.1016/0092-8674(77)90044-7).
- Marcus PI, Sekellick MJ, Nichol ST. 1992. Interferon induction by viruses. XXI. Vesicular stomatitis virus: interferon inducibility as a phylogenetic marker. *J Interferon Res* 12:297-305. <https://doi.org/10.1089/jir.1992.12.297>.
- Ferran MC, Lucas-Lenard JM. 1997. The vesicular stomatitis virus matrix protein inhibits transcription from the human beta interferon promoter. *J Virol* 71:371-377.
- Black BL, Rhodes RB, McKenzie M, Lyles DS. 1993. The role of vesicular stomatitis virus matrix protein in inhibition of host-directed gene expression is genetically separable from its function in virus assembly. *J Virol* 67:4814-4821.
- Petersen JM, Her L-S, Varvel V, Lund E, Dahlberg JE. 2000. The matrix protein of vesicular stomatitis virus inhibits nucleocytoplasmic transport when it is in the nucleus and associated with nuclear pore complexes. *Mol Cell Biol* 20:8590-8601. <https://doi.org/10.1128/MCB.20.22.8590-8601.2000>.
- Marcus PI, Sekellick MJ. 1994. Interferon induction: regulation by both virus and cell. *Hokkaido Igaku Zasshi* 69:1320-1331.
- Marcus PI, Sekellick MJ, Spiropoulou CF, Nichol ST. 1993. Interferon induction by viruses. XXII. Vesicular stomatitis virus-Indiana: M-protein and leader RNA do not regulate interferon induction in chicken embryo cells. *J Interferon Res* 13:413-418. <https://doi.org/10.1089/jir.1993.13.413>.
- Rodriguez LL, Pauszek SJ, Bunch TA, Schumann KR. 2002. Full-length genome analysis of natural isolates of vesicular stomatitis virus (Indiana 1 serotype) from North, Central and South America. *J Gen Virol* 83:2475-2483. <https://doi.org/10.1099/0022-1317-83-10-2475>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114-2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29:15-21. <https://doi.org/10.1093/bioinformatics/bts635>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455-477. <https://doi.org/10.1089/cmb.2012.0021>.
- Wheeler DL, Church DM, Federhen S, Lash AE, Madden TL, Pontius JU, Schuler GD, Schriml LM, Sequeira E, Tatusova TA, Wagner L. 2003. Database resources of the National Center for Biotechnology. *Nucleic Acids Res* 31:28-33. <https://doi.org/10.1093/nar/gkg033>.